

Excitation spectrum of a toroidal spin-1 Bose-Einstein condensate

H. Mäkelä^{1,2} and E. Lundh²

¹*QCD Labs, COMP Centre of Excellence, Department of Applied Physics,
Aalto University, P.O. Box 13500, FI-000 76 AALTO, Finland*

²*Department of Physics, Umeå University, SE-901 87 Umeå, Sweden*

We calculate analytically the excitation spectrum of a toroidal spin-1 Bose-Einstein condensate that is subjected to a homogeneous magnetic field and contains vortices with arbitrary winding numbers in the $m_F = \pm 1$ components of the hyperfine spin. The spectrum can be tuned by varying external parameters, such as the strength of the magnetic field. A rotonlike spectrum can be obtained, or an initially stable condensate can be made unstable by adjusting the magnitude of the magnetic field or trapping frequencies. The structure of the instabilities can be analyzed by measuring the particle density of the spin components.

PACS numbers: 03.75.Kk, 03.75.Mn, 67.85.De, 67.85.Fg

Bose-Einstein condensates (BECs) confined in toroidal traps have been subject to many experimental studies recently [1–6]. This research cover topics such as the observation of persistent current [1], phase slips across a stationary barrier [2], stochastic [3] and deterministic [4] phase slips between vortex states, the use of toroidal condensates in interferometry [5], and the stability of superfluid flow in a spinor condensate [6]. These experiments have given rise to theoretical studies discussing, e.g., the excitation spectrum and critical velocity of a superfluid BEC [7] and the simulation of the experiment [2] using the Gross-Pitaevskii equation [8, 9] and the truncated Wigner approximation [8]. Most of the experimental and theoretical studies concentrate on the properties of persistent currents. The phase of a toroidal BEC changes by $2\pi k$ as the toroid is encircled, the integer k being the winding number of the vortex. In a singly-connected geometry a vortex with $|k| > 1$ is typically unstable against splitting into vortices with smaller k . In a multiply connected geometry this process is suppressed for energetic reasons. In Ref. [3] it was shown experimentally that a vortex with winding number three can persist in a toroidal single-component BEC for up to a minute. In other words, toroidal geometry makes it possible to avoid the fast vortex splitting taking place in a singly-connected BEC and study the properties of vortices with large winding number.

In this letter, we calculate the linear excitation spectrum of a toroidal quasi one-dimensional (1D) spin-1 BEC. Motivated by the experimental results of Refs. [3, 6], we assume that the splitting of vortices occurs on a very long timescale in a spinor condensate where only one spin component is populated. The dominant instabilities can then be assumed to arise from the spin-spin interaction. For related theoretical studies on toroidal two-component condensates, see, for example, [10, 11]. In our analysis the population of the $m_F = 0$ component is taken to be zero initially, making it possible to calculate the excitation spectrum analytically. This type of state is straightforward to prepare experimentally and makes

it possible to observe the proliferation of instabilities by measuring the density of the $m_F = 0$ component.

The order parameter of a spin-1 Bose-Einstein condensate is $\psi = (\psi_1, \psi_0, \psi_{-1})^T$, where T denotes the transpose, and it fulfills the identity $\psi^\dagger \psi = n_{3D}$, where n_{3D} is the total particle density. The condensate is confined in a toroidal trap given in cylindrical coordinates as $U(r, z, \varphi) = m [\omega_r^2(R - r)^2 + \omega_z^2 z^2] / 2$, where R is the radius of the torus and ω_r, ω_z are the trapping frequencies in the radial and axial directions, respectively. We assume that the condensate is quasi 1D, so that the order parameter factors as $\psi(r, z, \varphi; t) = \psi_{r,z}(r, z) \psi_\varphi(\varphi; t)$, where $\psi_{r,z}$ is complex-valued and time independent. If the system is exposed to a homogeneous magnetic field oriented along the z -axis the energy functional becomes

$$E_{1D}[\psi_\varphi] = \int_0^{2\pi} d\varphi \left\{ \psi_\varphi^\dagger \left(-\epsilon \frac{\partial^2}{\partial \varphi^2} - \mu - p \hat{F}_z + q \hat{F}_z^2 \right) \psi_\varphi + \frac{1}{2} n \left[g_0 (\psi_\varphi^\dagger \psi_\varphi)^2 + g_2 \left(\psi_\varphi^\dagger \hat{\mathbf{F}} \psi_\varphi \right)^2 \right] \right\}, \quad (1)$$

where $\epsilon \equiv \hbar^2 / 2mR^2$, μ is an effective one-dimensional chemical potential, and $\hat{\mathbf{F}} = (\hat{F}_x, \hat{F}_y, \hat{F}_z)$ is the (dimensionless) spin operator of a spin-1 particle. The magnetic field introduces the linear and quadratic Zeeman terms, given by p and q , respectively. The sign of q can be controlled experimentally by using a linearly polarized microwave field [12]. The strength of the atom-atom interaction is characterized by $g_0 = 4\pi\hbar^2(a_0 + 2a_2)/3m$ and $g_2 = 4\pi\hbar^2(a_2 - a_0)/3m$, where a_F is the s -wave scattering length for two atoms colliding with total angular momentum F . The scattering lengths of ^{87}Rb used here are $a_0 = 101.8a_B$ and $a_2 = 100.4a_B$ [13], measured in units of the Bohr radius a_B . For ^{23}Na the corresponding values are $a_0 = 50.0a_B$ and $a_2 = 55.1a_B$ [14]. The magnetization in the z -direction, $f_z \equiv \int d\varphi \psi_\varphi^\dagger(\varphi) \hat{F}_z \psi_\varphi(\varphi) / N$, is a conserved quantity; the corresponding Lagrange multiplier can be included into p . Here N is the condensate particle number. We assume that in the initial state the spin is parallel to the z axis. In [15] it was argued that

in a homogeneous system the most unstable states are almost always of this form. The initial state reads

$$\psi_{\parallel}(\varphi) = \left(e^{ik_1\varphi} \sqrt{\frac{1+f_z}{2}}, 0, e^{i\theta} e^{ik_{-1}\varphi} \sqrt{\frac{1-f_z}{2}} \right)^T, \quad (2)$$

where θ is the relative phase and the integer $k_{\pm 1}$ is the winding number of the $m_F = \pm 1$ component. The energy and stability are independent of θ and therefore we set $\theta = 0$. If $k_1 = 1$ and $k_{-1} = 0$, $\psi_{\parallel}$ describes a half-quantum vortex (Alice string), see, e.g., Refs. [16–18]. The populations of $\psi_{\parallel}$ are time independent and the Hamiltonian reads $\hat{H}_{\parallel} = (g_0 n - \mu) \hat{I} + (g_2 n f_z - p_{\text{eff}} \hat{F}_z + q_{\text{eff}} \hat{F}_z^2)$, where $p_{\text{eff}} = p - \epsilon(k_1^2 - k_{-1}^2)/2$ and $q_{\text{eff}} = q + \epsilon(k_1^2 + k_{-1}^2)/2$. The time evolution operator of $\psi_{\parallel}$ is $\hat{U}_{\parallel}(t) = e^{-it\hat{H}_{\parallel}/\hbar}$. We calculate the excitation spectrum in a basis where $\psi_{\parallel}$ is stationary [15, 19]. In this basis, the energy of an arbitrary state ψ is given by $E_{\text{1D}}^{\text{new}}[\psi] \equiv E_{\text{1D}}[\hat{U}_{\parallel}\psi] + i\hbar\langle\psi| \left(\frac{\partial}{\partial t} \hat{U}_{\parallel}^{-1} \right) \hat{U}_{\parallel} \psi \rangle$ and the time evolution of ψ can be obtained from $i\hbar\partial\psi/\partial t = \delta E_{\text{1D}}^{\text{new}}[\psi]/\delta\psi^\dagger$. We write $\psi(\varphi; t) = \psi_{\parallel}(\varphi) + \delta\psi(\varphi; t)$ and expand the time evolution equation to first order in $\delta\psi$. The perturbation is written as

$$\delta\psi_j(\varphi; t) \equiv e^{ik_j\varphi} \sum_{s=0}^{\infty} u_{j;s}(t) e^{is\varphi} - v_{j;s}^*(t) e^{-is\varphi}, \quad (3)$$

where $j = 0, \pm 1$ and $k_0 \equiv 0$. Straightforward calculation gives the equation

$$i\hbar \frac{\partial}{\partial t} \begin{pmatrix} \mathbf{w}_0 \\ \mathbf{w}_{1;-1} \end{pmatrix} = (\hat{B}_2 \oplus \hat{B}_4) \begin{pmatrix} \mathbf{w}_0 \\ \mathbf{w}_{1;-1} \end{pmatrix}, \quad (4)$$

where $\oplus$ indicates the direct sum of matrices and the subscript i gives the dimension of $\hat{B}_i$. Only $\hat{B}_2$ depends on time. The vectors $\mathbf{w}_0$ and $\mathbf{w}_{1;-1}$ are defined as $\mathbf{w}_0 = (u_{0;s}, v_{0;s-k_1-k_{-1}})^T$ and $\mathbf{w}_{1;-1} = (u_{1;s}, u_{-1;s}, v_{1;s}, v_{-1;s})^T$. The matrix elements of $\hat{B}_2$ read

$$(\hat{B}_2)_{11} = \epsilon s^2 + g_2 n, \quad (5)$$

$$(\hat{B}_2)_{12} = -e^{-i[\epsilon(k_1^2 + k_{-1}^2) + 2q]t} g_2 n \sqrt{1 - f_z^2}, \quad (6)$$

$$(\hat{B}_2)_{22} = \epsilon(s - k_1 - k_{-1})^2 + g_2 n, \quad (7)$$

and $(\hat{B}_2)_{21} = -(\hat{B}_2)_{12}^*$. The operator $\hat{B}_4$ can be written as

$$\hat{B}_4 = \begin{pmatrix} \epsilon s^2 \hat{I}_2 + \hat{X} + \hat{D} & -\hat{X} \\ \hat{X} & -\epsilon s^2 \hat{I}_2 - \hat{X} + \hat{D} \end{pmatrix}, \quad (8)$$

where $\hat{I}_2$ is the 2×2 identity matrix, $\hat{D} = 2\epsilon s \text{diag}(k_1, k_{-1})$, and $\hat{X}$ is defined as

$$\hat{X}_{11,22} = 2g_+(1 \pm f_z), \quad \hat{X}_{12} = 2g_- \sqrt{1 - f_z^2}, \quad (9)$$

and $\hat{X}_{21} = \hat{X}_{12}$. Here $g_{\pm} \equiv (g_0 \pm g_2)n/4$ and $+$ ($-$) refers to $\hat{X}_{11}$ ($\hat{X}_{22}$). The 6×6 matrix $\hat{B}_2 \oplus \hat{B}_4$ is in

a block diagonal form, the blocks being the $\hat{B}_2$ and $\hat{B}_4$ matrices. It is then possible to calculate the eigenvalues and eigenvectors of $\hat{B}_2$ and $\hat{B}_4$ independently. $\hat{B}_2$ affects only the $m_F = 0$ component, that is, the magnetization remains unchanged but the direction of the spin vector is modified. The corresponding modes are therefore called spin modes. For similar reasons, the eigenvalues of $\hat{B}_4$ are called magnetization modes.

The eigenvalues of $\hat{B}_4$ can be calculated straightforwardly but they are too long to be shown here. The eigenvalues are independent of q and can be written as $\hbar\omega_j(s) = 2\epsilon s k_+ + \hbar\tilde{\omega}_j(s)$, where $\tilde{\omega}_j$ depends on $k_- \equiv (k_1 - k_{-1})/2$ but is independent of $k_+ \equiv (k_1 + k_{-1})/2$. Consequently, modes with differing k_+ but equal k_- have identical stability. We denote the real and imaginary parts of ω_j by ω_j^r and ω_j^i , respectively. Magnetization modes are unstable if at least one ω_j^i is positive. If $f_z = 0$ the eigenvalues simplify and are given by

$$\begin{aligned} \hbar\omega_{1,2,3,4}(s)|_{f_z=0} &= 2\epsilon s k_+ \\ &\pm 2\sqrt{\epsilon s^2 \left[\frac{\epsilon}{4}(s^2 + 4k_-^2) + g_+ \pm \sqrt{\epsilon k_-^2 (\epsilon s^2 + 4g_+) + g_-^2} \right]}. \end{aligned} \quad (10)$$

The signs are defined such that $++$, $-+$, $+-$, and $--$ correspond to $\omega_1, \omega_2, \omega_3$, and ω_4 , respectively. Unstable modes appear when the term inside the square brackets becomes negative. For rubidium and sodium $g_+ > 0$, guaranteeing that ω_1 and ω_2 are real. Only ω_3 can have a positive imaginary part, see Fig. 1. The amplitude




FIG. 1. The amplitudes of the unstable spin and magnetization modes for rubidium and sodium. Here $\epsilon = 0.75|g_2|n$, $q = 0$, and the unit of $\omega_{3,5}^i$ is $|g_2|n/\hbar$. The lines have been drawn by treating s as a continuous parameter; dots indicate the actual allowed non-vanishing values of $\omega_{3,5}^i$. In (c) and (d) the curves are reflection symmetric with respect to $s = k_+ = (k_1 + k_{-1})/2$.

of the unstable modes grows as $|k_-|$ increases. Now the allowed values of s are integers; the modes corresponding to $s = 0$ are always stable, but unstable modes are

present for $s = 1, 2, \dots, \lfloor \sqrt{4k_-^2 - 2g_2n/\epsilon} \rfloor$. Here $\lfloor \dots \rfloor$ indicates the floor function. It is therefore not possible to choose the parameters in such a way that a single mode $s > 1$ is unstable. A lower bound for the value of ϵ yielding at least one unstable mode in a sodium BEC is given by the equation $\epsilon(4k_-^2 - 1) \geq 2g_2n$. Because $\epsilon > 0$, the magnetization modes of a sodium condensate ($g_2 > 0$) with $k_- = 0$ and $|k_-| = 1/2$ are always stable. This is visualized in Fig. 1(b), where $\omega_3^i(s)$ corresponding to $(k_1, k_{-1}) = (0, 0)$ and $(k_1, k_{-1}) = (2, 1)$ is seen to vanish for every s . In a rubidium condensate ($g_2 < 0$) with $k_- = 0$ unstable modes exist if $\epsilon \leq 2|g_2|n$; if $|k_-| > 0$, instabilities are present regardless of the value of ϵ . For both rubidium and sodium the wavenumber of the fastest-growing instability is approximately given by the integer closest to $\sqrt{2/3} \sqrt{4k_-^2 - 2g_2n/\epsilon}$. Assume next that there is one dominant unstable mode. This is in particular the case if only the $s = 1$ mode is unstable. Using the eigenvector corresponding to $\omega_3^i(s)|_{f_z=0}$ we find that the position-dependent population difference $\Delta F_z(\varphi; t) \equiv \rho_1(\varphi; t) - \rho_{-1}(\varphi; t)$, where $\rho_j = |(\psi_j)_j + \delta\psi_j|^2$, is given by

$$\Delta F_z(\varphi; t) = \alpha e^{\omega_3^i t} \cos(\theta) + \beta e^{2\omega_3^i t} \sin(2\theta), \quad (11)$$

where $\theta = \arg[u_{1;s}(0)]/2 + s\varphi + \omega_3^i t$ and α and β are constants that depend on s, k_- , and the interaction coefficients. The presence of unstable magnetization modes can be experimentally confirmed by measuring $\Delta F_z(\varphi)$. If $k_- = 0$ we obtain $\beta = 0$, leading to ΔF_z having s peaks as a function of the coordinate φ . For nonzero k_- both α and β are in general nonvanishing and ΔF_z is a sum of two competing terms with periodicity s and $2s$. If $\omega_3^i(s) > 0$ we get $\omega_3^i(s) = 2\epsilon s k_+$, so that the nodes of ΔF_z rotate around the torus as time evolves if k_+ is nonzero. We see that although the value of k_+ does not affect the stability, it can alter the behavior of the density perturbations resulting from the instabilities.

We now turn to the spin modes. The time dependence of $\hat{B}_2$ can be eliminated by a simple change of basis. In the new basis, the eigenvalues giving the spin modes read

$$\hbar\omega_{5,6}(s) = 2\epsilon k_+(s - k_+) \quad (12)$$

$$\pm \sqrt{\{\epsilon[(s - k_+)^2 - k_-^2] + g_2n - q\}^2 - (1 - f_z^2)(g_2n)^2},$$

where $+$ ($-$) corresponds to ω_5 (ω_6). If $k_+ = 0$, the effect of vortices can be taken into account by scaling $q \rightarrow q_{\text{eff}} = q + \epsilon k_-^2$, i.e., the spin modes of a system with $(k_1, k_{-1}) = (k, -k)$ and $q = \tilde{q}$ are equal to the spin modes of a vortex-free condensate with $q = \tilde{q} + \epsilon k^2$. Spin modes are unstable if and only if the term inside the square root is negative. Now only ω_5 can have a positive imaginary part. The fastest-growing unstable mode is obtained at $\epsilon[(s - k_+)^2 - k_-^2] + g_2n - q = 0$ and has the amplitude $\hbar\omega_5^i(s) = |g_2|n\sqrt{1 - f_z^2}$. Unlike in the case

of the magnetization modes, the maximal amplitude is bounded from above and is independent of the winding numbers. By adjusting the strength of the magnetic field, the fastest-growing unstable mode can be chosen to be located at a specific value of s , showing that it is easy to adjust the stability properties experimentally. At $f_z = 0$ the width of the region on the s -axis giving positive ω_5^i is $|\sqrt{k_-^2 + q/\epsilon} - \sqrt{k_-^2 + q/\epsilon - 2g_2n/\epsilon}|$. This region can thus be made narrower by increasing ϵ, k_- , or q . Since the magnetization modes are insensitive to the magnetic field, the properties of the spin and magnetization modes can be tuned independently. The winding number dependence of unstable spin modes is illustrated in Figs. 1(c)-(d). Interestingly, by tuning ϵ and q a rotonlike spectrum




FIG. 2. The real (ω_5^r) and imaginary (ω_5^i) component of ω_5 (spin mode) for (a) rubidium and (b) sodium. Here $k_1 = -k_{-1}$, k_1 is an arbitrary integer, and $\epsilon = 0.3|g_2|n$. For the blue solid and blue dotted lines $q_{\text{eff}} = q + \epsilon k_-^2 = 5.5|g_2|n$ and for the orange dashed line $q_{\text{eff}} = -2|g_2|n$. The unit of $\omega_5^{r,i}$ is $|g_2|n/\hbar$. The blue solid and orange dashed lines show ω_5^r and the blue dotted line gives ω_5^i . These lines have been drawn by treating s as a continuous parameter; dots (circles) indicate the actual allowed non-vanishing values of ω_5^r (ω_5^i).

can be realized, see the solid and dotted blue lines in Fig. 2. Now the phonon part of the spectrum is missing, but the roton-maxon feature is present. For $f_z = k_+ = 0$ the roton spectrum exists if $q \geq \max\{0, 2g_2n\}$. Because only integer values of s are allowed, it may happen that the imaginary parts ω_3^i and ω_5^i are nonzero only in some interval of the s axis that does not contain integers, see Figs. 1(c)-1(d) and Fig. 2(a). In this case rotonic excitations are stable. Alternatively, there can be unstable modes close to the roton minimum, see Fig. 2(b) and Ref. [20]. As evidenced by the orange dashed lines in Fig. 2, the roton spectrum can be made to vanish simply by decreasing q . It is known that a rotonlike spectrum can exist in various types of BECs, such as in a dipolar condensate (see, e.g., [21–23]), in a Rydberg-excited condensate [24], or in a spin-1 sodium condensate prepared in a specific state [20]. In the present case the rotonlike spectrum exists both in a sodium and rubidium BEC and the state [Eq. (2)] giving rise to it is easy to prepare experimentally. Note that the roton-maxon feature exists also in a vortex-free condensate and for any $|f_z| < 1$. These results suggest that the roton-maxon character of the spectrum is rather a rule than an exception in spinor BECs. By changing the value of q an unstable spin mode

can be introduced or made to vanish. For example, using the parameter values corresponding to the blue solid line in Fig. 2(a), we find that by decreasing (increasing) the value of q_{eff} from $5.5|g_2|n$ to $4|g_2|n$ ($6|g_2|n$), the $s = 4$ ($s = 5$) modes can be made unstable. This opens the way for quench experiments of the type described in [25, 26]. In the present system the number of excited unstable modes and their wavenumbers can be controlled with a high precision. Instead of altering q , instabilities can also be induced by making ϵ smaller by changing the trapping frequencies.

The properties of unstable spin modes can be studied experimentally by measuring the density and phase of the $m_F = 0$ spin component. Assuming that there is one dominant unstable spin mode located at wavenumber s , we find the density of the $m_F = 0$ component to be

$$\rho_0(\varphi; t) = |\delta\psi_0(\varphi; t)|^2 \approx e^{2\omega_5^i t} \left\{ |u|^2 + |v|^2 - 2|uv| \cos[\theta_u + \theta_v + 2\omega_5^r t + 2(s - k_+) \varphi] \right\}, \quad (13)$$

where $u = u_{0,s}(0)$, $v = v_{0,s-k_1-k_{-1}}(0)$, θ_u is defined through $u = |u|e^{i\theta_u}$, and θ_v is defined similarly. The number of nodes in ρ_0 is $2|s - k_+|$, that is, if $2k_+$ is even (odd), only an even (odd) number of nodes can exist. The density peaks resulting from an instability rotate around the torus if $\omega_5^r = 2\epsilon k_+(2s - k_+)$ is nonzero. The ratio of the components of the eigenvector $\mathbf{w}_0 = (u, v)^T$ corresponding to ω_5 reads $v/u = (a - b)/\sqrt{1 - f_z^2 g_2 n}$, where $a = \epsilon[(s - k_+)^2 - k_{-1}^2] + g_2 n - q$ and $b = \sqrt{a^2 - (1 - f_z^2)(g_2 n)^2}$. If there is a dominant unstable mode with wavenumber s we get $|v/u| = 1$ and Eq. (13) gives $\rho_0(\varphi; t) \approx 4|u|^2 e^{2\omega_5^i t} \sin^2[\omega_5^r t + (s - k_+) \varphi]$, where we have set $\theta_u = \theta_v = 0$ for simplicity. The density vanishes at $\varphi = (\omega_5^r t + n\pi)/(k_+ - s)$, where n is an arbitrary integer. From the fact that $\delta\psi_0 \sim \sin[\omega_5^r t + (s - k_+) \varphi]$ we see that the sign of $\delta\psi_0$ changes at every point where the density vanishes. This holds true regardless of the value of f_z . The $\pm\pi$ phase jump at the density minima resembles the behavior of the phase of a dark soliton [27].

We look next at the values of ϵ corresponding to two recent experiments. Using the Thomas-Fermi (TF) approximation and the parameter values of the sodium experiment [2] we get $\epsilon \approx 0.04g_2n$. This means that magnetization modes are stable if $|k_-| \leq 3$, but spin modes are unstable even when $|k_-| = 0$. In the following k_- is such that only spin modes are unstable. If $0 < q \leq 0.04g_2n$, $f_z = 0$, and $(k_1, k_{-1}) = (0, 0)$, the instability appears as a position-independent, homogeneous, increase in ρ_0 . On the other hand, if $(k_1, k_{-1}) = (1, 0)$ we get $\rho_0(\varphi; t) \sim e^{2\omega_5^i t} \sin^2[(\epsilon t + \varphi)/2]$, so that the density ρ_0 is peaked around $\varphi = -\epsilon t + \pi$. This illustrates that even a small ϵ can lead to a strongly winding number dependent density ρ_0 . The parameters of the rubidium experiment [6] give $\epsilon \approx 0.18|g_2|n$. Now some magnetization modes are unstable regardless of the values of winding numbers.

If $q_{\text{eff}} \approx 2|g_2|n$ the $s = 4$ spin mode, situated close to the minimum of the roton spectrum, is the only unstable spin mode. At $q_{\text{eff}} = 0$, the $s = 1, 2, 3$ modes are unstable, and at $q_{\text{eff}} = -2|g_2|n$ all spin modes are stable. Using the TF approximation we get $\epsilon \sim (\omega_r \omega_z N R^3)^{-1/2}$, showing that the stability of the magnetization modes can be controlled by adjusting the trapping frequencies, number of particles, and the radius of the toroid.

The first experimental realization of a toroidal spin-1 BEC was reported very recently [6]. The stability of a rubidium BEC with vortices with equal winding numbers in the $m_F = 1$ and $m_F = 0$ components was found to depend strongly on the population difference of the two components, the most unstable situation corresponding to equal population. Although not directly comparable, our analysis agrees qualitatively with this result: The amplitude of unstable spin and magnetization modes increases as $f_z \rightarrow 0$.

In conclusion, we have calculated analytically the linear excitation spectrum of a toroidal spin-1 BEC. In this calculation the strength of the magnetic field and the winding numbers of the $m_F = \pm 1$ components were treated as free parameters. We found that the shape of the spectrum can be tuned by adjusting the strength of the magnetic field. In particular, a spectrum with a roton-maxon structure can be realized both in rubidium and sodium condensates. Furthermore, by varying the magnetic field, an initially stable condensate can be made unstable. We have also shown that some unstable modes lead to a dark soliton-like wavefunction of the $m_F = 0$ spin component. Finally, by considering two recent experiments on toroidal single-component BECs, we argued that various types of instabilities can be realized in a spin-1 condensate having the same parameter values as these experiments.

This research has been supported by the Alfred Kordelin Foundation and the Academy of Finland through its Centres of Excellence Program (Project No. 251748).

-
- [1] C. Ryu, M. F. Andersen, P. Cladé, V. Natarajan, K. Helmerson, and W. D. Phillips, Phys. Rev. Lett. **99**, 260401 (2007).
 - [2] A. Ramanathan, K. C. Wright, S. R. Muniz, M. Zelán, W. T. Hill III, C. J. Lobb, K. Helmerson, W. D. Phillips, and G. K. Campbell, Phys. Rev. Lett. **106**, 130401 (2011).
 - [3] S. Moulder, S. Beattie, R. P. Smith, N. Tammuz, and Z. Hadzibabic, Phys. Rev. A **86**, 013629 (2012).
 - [4] K.C. Wright, R.B. Blakestad, C.J. Lobb, W.D. Phillips, and G.K. Campbell, Phys. Rev. Lett. **110**, 025302 (2013).
 - [5] G.E. Marti, R. Olf, and D.M. Stamper-Kurn, arXiv:1210.0033.
 - [6] S. Beattie, S. Moulder, R.J. Fletcher, and Z. Hadzibabic, Phys. Rev. Lett. **110**, 025301 (2013).

- [7] R. Dubessy, T. Liennard, P. Pedri, and H. Perrin, Phys. Rev. A **86**, 011602(R).
- [8] A. C. Mathey, C. W. Clark, and L. Mathey, arXiv:1207.0501.
- [9] F. Piazza, L. A. Collins, and A. Smerzi, arXiv:1208.0734.
- [10] J. Smyrnakis, S. Bargi, G.M. Kavoulakis, M. Magiropoulos, K. Kärkkäinen, and S.M. Reimann, Phys. Rev. Lett. **103**, 100404 (2009).
- [11] K. Anoshkin, Z. Wu, and E. Zaremba, arXiv:1207.3449.
- [12] F. Gerbier, A. Widera, S. Fölling, O. Mandel, and I. Bloch, Phys. Rev. A **73**, 041602(R) (2006).
- [13] E. G. M. van Kempen, S. J. J. M. F. Kokkelmans, D. J. Heinzen, and B. J. Verhaar, Phys. Rev. Lett. **88**, 093201 (2002).
- [14] A. Crubellier, O. Dulieu, F. Masnou-Seeuws, M. Elbs, H. Knöckel, and E. Tiemann, Eur. Phys. J. D. **6**, 211 (1999).
- [15] H. Mäkelä, M. Johansson, M. Zelan, and E. Lundh, Phys. Rev. A **84**, 043646 (2011).
- [16] U. Leonhardt and G.E. Volovik, JETP Lett. **72**, 66 (2000).
- [17] T. Isoshima, K. Machida, and T. Ohmi, J. Phys. Soc. Jpn. **70**, 1604 (2001).
- [18] S. Hoshi and H. Saito, Phys. Rev. A **78**, 053618 (2008).
- [19] H. Mäkelä and E. Lundh, Phys. Rev. A **85**, 053622 (2012).
- [20] M. Matuszewski, Phys. Rev. Lett. **105**, 020405 (2010).
- [21] D.H.J. O'Dell, S. Giovanazzi, and G. Kurizki, Phys. Rev. Lett. **90**, 110402 (2003).
- [22] L. Santos, G.V. Shlyapnikov, and M. Lewenstein, Phys. Rev. Lett. **90**, 250403 (2003).
- [23] R.W. Cherno and E. Demler, Phys. Rev. Lett. **103**, 185301 (2009).
- [24] N. Henkel, R. Nath, and T. Pohl, Phys. Rev. Lett. **104**, 195302 (2010).
- [25] L. E. Sadler, J. M. Higbie, S. R. Leslie, M. Vengalattore, and D. M. Stamper-Kurn, Nature **443**, 312 (2006).
- [26] E.M. Bookjans, A. Vinit, and C. Raman, Phys. Rev. Lett. **107**, 195306 (2011).
- [27] D.J. Frantzeskakis, J. Phys. A **43**, 213001 (2010).